

Alasdair P. MacGowan

Brinell Centre for Antimicrobial Research and Evaluation, Southmead Health Services, NHS Trust; and University of Bristol, Department of Medical Microbiology, Southmead Hospital, Wootton Bassett, Wiltshire, United Kingdom

Summary: The glycopeptide antibacterial drugs, vancomycin and teicoplanin, are widely used in hospitals for therapy of severe or multibacterial infections that has a positive result on Gram's stain test. Although vancomycin resistance is common in some hospital-acquired *Enterococcus* sp and resistance to teicoplanin occurs among *Staphylococci* sp glycopeptides remain the cornerstone of therapy for infection due to methicillin-resistant *Staphylococcal aureus* (MRSA), coagulase-negative *Staphylococci* organisms, and infection related to implanted devices. Therapeutic drug monitoring (TDM) of these agents remains controversial, but advances in our understanding of their pharmacodynamics and further clinical studies are helping clarify the situation. In the future, a more rational approach to monitoring will probably result in less intensive monitoring of vancomycin but more intensive monitoring of teicoplanin. **Key Words:** Vancomycin—Teicoplanin—Pharmacodynamics—Therapeutic drug monitoring.

Postantibiotic effect is the ability of an antimicrobial to suppress the generation of bacteria for several hours after antimicrobial concentrations have fallen below the MIC. Although the exact mechanism of the PAE is unknown, it may be related to repair of damaged, but not killed, cells; separation of bound drug from target; or synthesis of new enzymes or proteins (1). Penicillins, cephalosporins, macrolides, and aminoglycosides have a PAE against bacteria that have a positive result on Gram's stain tests (2). Although the measurement of the PAE depends on the method used (3), vancomycin has been

Received February 11, 1998; accepted July 23, 1998.
Address correspondence and reprint requests to Alan H. P.
Mantovani, University of Bristol, Department of Medical Microbiol-
ogy, St. Mark's Hospital, Westbury-on-Trym, Bristol BS10 5DB,
United Kingdom.

Telaprevir also has a PAE that appears to be longer than that of vancomycin (8,7). If a concentration well below the MIC is allowed to remain instead of completely removing the antibiotic during the measurement of the PAE, the PAE duration is doubled for vancomycin, which is termed a sub-MIC effect. This measure is akin to the exponential decay of a drug concentration in serum (8).

Increasing the concentration of vancomycin in the therapeutic range (i.e., from 5 mg/L to 40 mg/L) does not increase the time to kill 99.9% of the bacterial population or the rate of kill (9); the rate of killing is slower for teicoplanin than for vancomycin, perhaps because of the former's high protein binding (10).

The pharmacodynamics of glycopeptides studied in several animal models support the concept that high initial concentrations offer no advantage in bacterial killing or mortality, whereas higher, sustained concentrations or more frequent dosing have improved survival in animal models of infective endocarditis (11,12).

is a complex analysis using a more model, multiple

pharmacodynamic parameters were compared with the effective dose $_{50}$ (ED_{50}). The time serum concentrations (T) exceeded the MIC ($T > MIC$) and was best related to ED_{50} when treating penicillin-resistant *Staphylococci* organisms with either vancomycin or teicoplanin (13). Use of an in vitro, continuous bacterial culture model supports this finding. By simulating four different therapeutic regimens with various peak or trough concentrations or areas under the curve (AUC), but maintaining $T > MIC$ at 100%, it was shown that there were no differences in the degree or rate of *S. aureus* killing (14).

Laboratory and animal evaluation of glycopeptide pharmacodynamics indicates that glycopeptides do not show concentration-dependent killing in the therapeutic range; hence, high peak concentrations are unlikely to be of benefit. In addition, they have a PAE and sub-MIC effects, indicating that serum concentrations need not exceed the MIC for all of the dosing interval and that $T > MIC$ or sustained concentrations are related to outcome. Protein binding affects bacterial killing with teicoplanin. Therefore, the dosing interval is probably best optimized as $T > MIC$ plus PAE, although clinically, with conventional doses of vancomycin, $T > MIC$ is 100%. This result has led some to propose that smaller doses than the standard 2 grams per day of vancomycin may be just as effective in clinical practice; alternatively, longer dosing intervals may be appropriate for glycopeptides (15).

PHARMACOKINETICS

The pharmacokinetics of vancomycin and teicoplanin have been extensively studied and are known to vary in different patient groups. For example, vancomycin handling is changed in renal impairment (16-18); obesity (19,28); liver failure (21); various renal support therapies (22-27); neutropenia (28); malignancy (29); age, and gender (30); and with sepsis and in therapy (Table 1 (31,32)).

Similarly, teicoplanin pharmacokinetics are altered in renal impairment (33,34), renal support therapies (35,36); children and the elderly (37); intravenous (IV) drug shunters (38); burn patients (9); and neutropenia (see Table 1 (40)). In addition, it is clear that standard dosing of teicoplanin (400 mg \times 2 for 24 hours, then 400 mg 24 hourly) results in significant numbers of patients having predose serum concentrations of 10 mg/L (40).

However, pharmacokinetic variability on its own can rarely be a justification for TDM and only becomes important if serum concentrations can be linked to toxicity or efficacy. This link has been tenuous for glycopeptides and continued TDM was questioned in the late 1980s and

TABLE 1. Patient factors affecting vancomycin or teicoplanin pharmacokinetics

Patient Factor	Pharmacokinetic change	Reference no.
Vancomycin		
Renal impairment	Increasing $t_{1/2}$ with decreasing creatinine clearance	17
Chronic intermittent hemodialysis	As for renal impairment: little drug removed by dialysis	22
Chronic intermittent peritoneal dialysis	Prolonged $t_{1/2}$	23
Continuous veno-venous hemofiltration or dialysis	Increased clearance compared to hemodialysis	21,27
Obesity	Shorter $t_{1/2}$, larger volume of distribution	18,19
Liver failure	Longer $t_{1/2}$	21
Age	Longer $t_{1/2}$ in neonates than infants and children	29,36
Sepsis	Prolonged $t_{1/2}$	25
Teicoplanin		
Renal impairment	Increasing $t_{1/2}$ with decline in creatinine clearance	33,34
Chronic intermittent hemodialysis	Reduced clearance	35
Chronic ambulatory peritoneal dialysis	Prolonged $t_{1/2}$, increased volume of distribution	32
Continuous veno-venous hemofiltration	Prolonged $t_{1/2}$	33
Continuous hemofiltration	$t_{1/2}$ not prolonged	36
Intermittent drug shunters	Increased clearance	38
Burn patients	Increased $t_{1/2}$	39
Neutropenia	Increased elimination, larger interindividual variability	40

early 1990s (42-46). This process has now resulted in the emergence of new data.

TOXICITY

Vancomycin serum monitoring, if performed, is aimed at reducing the risks of nephrotoxicity or ototoxicity; it will not reduce immediate or infusion-related toxicities. Ototoxicity is difficult to assess clinically and data is sketchy because they are often composed of case reports in patients with renal failure, who sometimes have high serum concentrations. It is not sufficient to make any association between concentrations and toxicity.

The incidence of nephrotoxicity is probably less than 5% in patients treated with vancomycin alone, but higher if a combination of vancomycin plus an aminoglycoside is used (47). Toxicity is also associated with longer courses of therapy and the original report of Faber and Moellering (47) linked three patients to trough concentrations of 30 mg/L to 65 mg/L before the onset of tox-

ically. There are now several reports that vancomycin TDM services involving prescription review, blood concentration measurement, and dose modification by clinical pharmacists can reduce the incidence of nephrotoxicity. In a 1994 prospective cohort study in a teaching hospital, 116 patients who received more than 4 days' therapy and were not neutropenic or in an intensive care unit or established renal failure were studied. It was shown that the rate of nephrotoxicity was 24% in patients not randomized to the TDM service, compared with 7% of those who received TDM (43). A similar and prospective randomized study in 70 patients with hematologic malignancy indicated that nephrotoxicity was lower in patients recruited into the TDM arm (mild toxicity, 13.5%; moderate, 0%) compared with those who received no TDM (mild, 33%; moderate, 9.1%) (49). Furthermore, in a retrospective review of 273 patients with positive results of infection with *Cran's* stain, it was shown that serum vancomycin concentrations before onset of nephrotoxicity were higher (23.5 [2.5 mg/L]) in those in whom toxicity developed than in those in whom it did not (10.2 [3.8 mg/L]) (50). In contrast, in a prospective study of patients randomized to have dose adjustment to achieve predose concentrations in the ranges of 5 mg/L to 10 mg/L, 10 mg/L to 15 mg/L, or 15 mg/L to 25 mg/L, no correlation was found to nephrotoxicity (51).

Thrombocytopenia associated with large doses of teicoplanin (30 mg/kg per day) has recently been related to trough concentrations; for those with trough concentrations of more than 60 mg/L, eight of 58 patients had a decrease in platelets, whereas with trough concentrations of less than 60 mg/L, 12 of 251 had a decrease in platelets ($p < 0.05$) (52).

OUTCOME

Retrospective data reviews of teicoplanin clinical trials have indicated that serum concentrations are related to clinical outcomes. In an open multicenter study in which most patients had right-sided infection due to *S. aureus* and predose concentrations had been adjusted to between 10 mg/L to 15 mg/L, it was reported that predose concentrations of teicoplanin of more than 40 mg/L were associated with improved outcomes (53). A further study (majority of bone infection due to *S. aureus*) suggested that larger doses than were conventionally used at that time were required for successful therapy and that average troughs were 36.3 ($n = 10$) in those successfully treated, and only 9.7 mg/L in the three clinical failures (54). A retrospective review of three trials in the United States indicated an association between increased dose, high trough serum levels, trough concentration/MIC ratio

and days to clear bacteremia, fewer days, and clinical improvement (55). A further retrospective review of 58 cases published with sufficient pharmacokinetic and susceptibility data for analysis indicated a relationship between predose and postdose teicoplanin concentrations and predose/MIC or postdose/MIC ratios and clinical outcomes. Dose was not related to outcome in this review, in which most of the patients had severe staphylococcal infection (56). In a more recent review of 92 patients with *S. aureus* bacteremia using a multivariate analysis to relate age, weight, dose, timing dose, combination therapy, and serum concentrations to outcome has shown that only trough concentration and age were significantly related to outcome (57). A prospective study of teicoplanin to treat *S. aureus* infective endocarditis showed that if the predose concentration was less than 20 mg/L, six of 10 patients failed, compared with one of 11 if the concentration was more than 20 mg/L ($p < 0.05$) (58).

The data relating to vancomycin serum concentration to efficacy is less clear. Two prospective studies showed that therapeutic drug monitoring (TDM) services had no effect on efficacy (48,49) and an intervention in which patients were deliberately stratified into three groups with predose targets of 5 mg/L to 10 mg/L, 10 mg/L to 15 mg/L, or 15 mg/L to 25 mg/L showed no difference in fever days or clinical outcome (51). In contrast, two retrospective reviews were able to relate serum concentrations to outcome measures. In a retrospective review of 273 patients with positive results of infection proven with *Cran's* stain, Zimolman and colleagues (50) were able to relate troughs of more than 10 mg/L to a reduced number of fever days and an improved white blood cell response but not to lengths of stay or mortality. Mulhern and colleagues (59) related trough concentrations to relapse rates in patients with peritonitis who were treated with continuous ambulatory peritoneal dialysis for end-stage renal disease. When the mean predose concentration was less than 12 mg/L, 9 of 16 patients relapsed; when it was more than 12 mg/L, none of 17 relapsed.

In conclusion, pharmacodynamic principles indicate that predose glycopeptides should be related to the outcome of infection measures. Evidence now exists in humans based on teicoplanin therapy of staphylococcal infections; the evidence is less conclusive for vancomycin.

For both vancomycin and teicoplanin, there is data to link predose concentrations to toxicity (nephrotoxicity for vancomycin and thrombocytopenia for teicoplanin). Table 2 summarizes present recommendations for glycopeptide TDM, including those which have been used and criticized in the past, and more streamlined recommendations that may be more appropriate for the future.

Pharm Drug Monit, Vol. 26, No. 3, 2004

TABLE 2. Recommendations for therapeutic drug monitoring glycopeptides

	Patient group	Target concentrations
Vancomycin	All patients on therapy >4 to 5 days (≥3 days if conservative)	Peak: 15-20 mg/L; Pre-dose: 5-10 mg/L; trough: 20-40 mg/L, if conservative
	Severe infection	Peak: >10 mg/L; Pre-dose: >10 mg/L (≥20 mg/L if conservative)
Telaprevir	Severe infection	Peak: >10 mg/L; Pre-dose: >10 mg/L (≥20 mg/L if conservative)
	Severe infection	Peak: >10 mg/L; Pre-dose: >10 mg/L (≥20 mg/L if conservative)
IVDA	Severe infection	Peak: >10 mg/L; Pre-dose: >10 mg/L (≥20 mg/L if conservative)
	Severe infection	Peak: >10 mg/L; Pre-dose: >10 mg/L (≥20 mg/L if conservative)

IE: Infective endocarditis; IVDA: Intravenous drug abuse.

REFERENCES

1. Gould IM. Pharmacokinetics and the relationship between in vivo and in vitro activity of antimicrobial agents. *J Chemother* 1977;2:4-13.
2. Sandhu RW, Gorbett AJ, Bohn DE, et al. Post antibiotic suppression of bacterial growth. *Rev Infect Dis* 1981;3:28-37.
3. MacKinnon PM, Gould IM. The post antibiotic effect. *J Antimicrob Chemother* 1983;12:519-57.
4. Hargreaves M, Nelson LS, Mihara M, et al. Post antibiotic effect of beta lactams on Gram-negative bacteria in relation to pharmacokinetics, initial killing and MIC. *Eur J Clin Microbiol Infect Dis* 1991;10:527-34.
5. Hargreaves M. Pharmacodynamic effects of antibiotics. Studies on bacterial morphology, initial killing, post antibiotic effect and effective regimens. *Stand J Infect Dis* 1992;11:3-32.
6. Teraoka K, Kouchi K, Shimizu K, et al. Post antibiotic effect and clinical evaluation of telaprevir. *Myken Kagaku Ryojin Gokai Kagaku* 1993;41:177-7.
7. Chappin MA, Yin YP, Ashby AP, et al. In vitro comparison of the post-antibiotic effect of vancomycin and telaprevir. *J Antimicrob Chemother* 1996;38:205-13.
8. Chabot-Torres J, Lavoie F, Chen Q. Post antibiotic sub-MIC effects of vancomycin, teicoplanin, rifampicin and rifaximin. *Antimicrob Agents Chemother* 1992;36:1824-8.
9. Lavoie F, Walker RJ, Roberts JM, et al. The concentration-independent effect of monosaccharide and disaccharide dyes in vancomycin concentrations on the killing of *Staphylococcus aureus* under aerobic and anaerobic conditions. *J Antimicrob Chemother* 1996;38:189-97.
10. Bailey EM, Rybak MJ, Kohn GW. Comparative effect of protein binding on the killing activities of telaprevir and vancomycin. *Antimicrob Agents Chemother* 1991;35:1070-72.
11. Chappin MA, Yin YP, Ashby AP, et al. The pharmacokinetics and comparative effect of telaprevir in rabbits and comparative efficacy with vancomycin in an experimental endocarditis model. *J Antimicrob Chemother* 1992;30:11-11.
12. Chabot-Torres J, Kennedy E. Effect of dosage, peak and trough concentrations in serum, protein binding and bactericidal rate on efficacy of telaprevir in a rabbit model with endocarditis. *Antimicrob Agents Chemother* 1992;36:1810-4.
13. Kohn GW, Roberts JM, Roberts JM, et al. Activity of vancomycin and telaprevir against penicillin-resistant pneumococci in vitro and in vivo and correlation to pharmacokinetic parameters in the mouse peritonitis model. *Antimicrob Agents Chemother* 1997;41:1919-21.
14. Duffell SB, Begg EJ, Chabot-Torres J, et al. Efficacy of different vancomycin dosing regimens against *Staphylococcus aureus* demonstrated with dynamic in vitro models. *Antimicrob Agents Chemother* 1994;38:2480-2.
15. McGowan AP, Bowler KE. Pharmacokinetics of antimicrobial agents and rationale for their dosing. *J Chemother* 1987;9:64-73.
16. MacKinnon PM, Knapton DL, Greenhill DJ. Pharmacokinetics of vancomycin in normal subjects and in patients with reduced renal function. *Rev Infect Dis* 1981;3:720-3.
17. Morris JR, McCarty RW, Hulseworth CF, et al. Pharmacokinetics of vancomycin in patients with various degrees of renal function. *Antimicrob Agents Chemother* 1984;23:613-7.
18. Oake BA, Quinton R, Digby EM, et al. Pharmacokinetics of vancomycin in adults. *Rev Infect Dis* 1981;3:720-3.
19. Blum RA, Bauer LA, Miller DD, et al. Vancomycin pharmacokinetics in normal and morbidly obese subjects. *Antimicrob Agents Chemother* 1982;21:573-82.
20. Vane-Bryne K, Gray DFP, Gifford SK, et al. Effect of obesity on vancomycin pharmacokinetic parameters as determined by using a Bayesian forecasting technique. *Antimicrob Agents Chemother* 1993;37:436-40.
21. Brown M, Ho DHW, Fong KLL, et al. Effect of hepatic function on vancomycin clinical pharmacology. *Antimicrob Agents Chemother* 1981;23:680-4.
22. Morris JR, Aronoff AS, Rasmussen BJ, et al. Vancomycin pharmacokinetics in patients with peritonitis on peritoneal dialysis. *Antimicrob Agents Chemother* 1982;22:710-4.
23. Blum RA, Hulseworth CF, Salas MCL, et al. Pharmacokinetics of vancomycin in patients undergoing continuous ambulatory peritoneal dialysis. *Antimicrob Agents Chemother* 1984;23:610-6.
24. Touraine MA, Patel RV, Anand JV, et al. Vancomycin removal by high-flux polyacrylonitrile membranes in critically ill patients with end-stage renal disease. *Am J Kidney Dis* 1993;21:669-74.
25. Gremelle F, Krawinkel M, de la Couraye H, et al. Clinical pharmacokinetics during continuous haemofiltration. *Clin Pharmacol Ther* 1994;56:177-71.
26. Albrecht T, Meier TA, et al. Comparison of the effects of three haemodialysis membranes on vancomycin disposition. *Int J Artif Organs* 1994;17:223-8.
27. Jones CL, Levy G, Singer M, et al. Pharmacokinetics of vancomycin during continuous haemofiltration. *Statist Case Med* 1993;19:347-50.
28. Le Normand Y, Milpied N, Kergueris MF, et al. Pharmacokinetic parameters of vancomycin for therapeutic regimens in end-stage renal patients. *Int J Biomed Comput* 1994;36:123-3.
29. Chang D. Influence of malnutrition on the pharmacokinetics of vancomycin in infants and children. *Pediatr Infect Dis J* 1993;14:647-73.
30. Saly RL, Brundage RC, Jacobs PD, et al. Population pharmacokinetics of vancomycin in neonates. *Clin Pharmacol Ther* 1994;56:169-73.
31. Rotstein JC, Cronin K, Zech D, et al. Pharmacokinetics of vancomycin in 28 patients and dosage recommendations. *Antimicrob Agents Chemother* 1982;22:391-4.
32. Pan L, Roubil M, Lepin E, et al. Changes in vancomycin pharmacokinetics during treatment. *Ther Drug Monit* 1986;8:149-53.
33. Strickland JA, Thompson GA, Heary MT, et al. Application of telaprevir dosage adjustment guidelines for orally impaired patients over the range 3 to 30 mg/L. *Therapeutic Drug Monit* 1992;14:371-81.
34. Derbyshire N, Webb DS, Roberts D, et al. Pharmacokinetics of telaprevir in subjects with varying degrees of renal function. *J Antimicrob Chemother* 1989;23:669-73.
35. Hoffer DM, Joseph P, Shumway R, et al. Pharmacokinetics of telaprevir haemodialysis patients. *Antimicrob Agents Chemother* 1991;35:204-7.
36. Kishimoto S, Peckow M, Leblond R, et al. Influence of ureteric dialysis on telaprevir elimination. *Br J Clin Pharmacol* 1990;50:75-7.

Ther Drug Monit, Vol 26, No 4, 1999

Supplied by the British Library - "The world's knowledge" www.bl.uk